arXiv:alg-geom/9312006v1 14 Dec 1993

AN ALGORITHMIC CRITERION FOR BASICNESS IN DIMENSION 2

Francesca Acquistapace*
Fabrizio Broglia*
Dipartimento di Matematica,
Università di Pisa, I56100 Pisa, Italy

M. Pilar Vélez†
Departamento de Geometría y Topología, Facultad de Matemáticas,
Universidad Complutense de Madrid, 28040 Madrid, Spain

December,13,1993

Mathematics Subject Classification: 14P20

Introduction.

In this paper we give a constructive procedure to check basicness of open (or closed) semialgebraic sets in a compact, non singular, real algebraic surface X. It is rather clear that if a semialgebraic set S can be separated from each connected component of $X \setminus (S \cup \partial_{\rm z} S)$ (when $\partial_{\rm z} S$ stands for the Zariski closure of $(\overline{S} \setminus \mathrm{Int}(S)) \cap \mathrm{Reg}(X)$), then S is basic. This leads to associate to S a finite family of sign distributions on $X \setminus \partial_{\rm z} S$; we prove the equivalence between basicness and two properties of these distributions, which can be tested by an algorithm described in [ABF].

By this method we find for surfaces a general result of [AR2] about the "ubiquity of Lojasiewicz's example" of non basic semialgebraic sets (2.9). There is a close relation between these two properties and the behaviour of fans in the algebraic functions field

---

*Members of CNSAGA of CNR, partially supported by MURST

†Partially supported by DGICYT and a Beca-Complutense

of X associated to a real prime divisor (Lemmas 3.5 and 3.6). We use this fact to get an easy proof (Theorem 3.8), for a general surface X, of the well known 4-elements fan's criterion for basicness (see [Br] and [AR1]). Furthermore, if the criterion fails, using the description of fans in dimension 2 [Vz], we find an algorithmic method to exhibit the failure. Finally, exploiting this thecnics of sign distribution we give one improvement of the 4-elements fan's criterion in [Br] to check if a semialgebraic set is principal.

In fact, one goal of this paper is to give purely geometric proofs, in the case of surfaces, of the theory of fans currently used in Semialgebraic Geometry ([Br]). In particular, we only need the definitions of that theory. It is at least remarkable that while the notion of fan is highly geometric in nature all known proofs of the main results are pure quadratic forms theory.

1 Geometric review of basicness

Let $X \subset \mathbb{R}^n$ be an algebraic surface. Denote by $\mathcal{R}(X)$ the ring of regular functions on X. Let $S \subset X$ be a *semialgebraic set*, that is

$$S = \bigcup_{i=1}^{p} \{x \in X : f_{i1}(x) > 0, \dots, f_{ir_i}(x) > 0, g_i(x) = 0\}$$

with $f_{i1}, \dots, f_{ir_i}, g \in \mathcal{R}(X)$, for $i = 1, \dots, p$.
We will simply write: $S = \bigcup \{f_{i1} > 0, \dots, f_{ir_i} > 0, g_i = 0\}$.

Definition 1.1 *A semialgebraic set S is* basic open *(resp.* basic closed*) if there exist $f_1, \dots, f_r \in \mathcal{R}(X)$ such that*

$$S = \{x \in X : f_1(x) > 0, \dots, f_r(x) > 0\}$$

$$(\textit{resp. } S = \{x \in X : f_1(x) \geq 0, \dots, f_r(x) \geq 0\})$$

Definition 1.2 *A semialgebraic set $S \subset X$ is* generically basic *if there exist a Zariski closed set $C \subset X$, with $\dim(C) \leq 1$, such that $S \setminus C$ is basic open.*

Denote by $\partial_z S$ the Zariski closure of the set $\partial(S) = (\overline{S} \setminus \mathrm{Int}(S)) \cap \mathrm{Reg}(X)$. It is known that in dimension 2 *basic* and *generically basic* have almost the same meaning (see [Br]). We give here a direct proof of this fact.

Lemma 1.3 *Let S be an open semialgebraic set in X, if S is generically basic then $S \cap \partial_z S$ is a finite set.*

Proof: Suppose there exist $f_1, \ldots, f_s \in \mathcal{R}(X)$ and an algebraic set $C \subset X$ such that $\dim(C) \leq 1$ and

$$S \setminus C = \{f_1 > 0, \ldots, f_s > 0\}.$$

Suppose also that there is an irreducible component H of $\partial_z S$ such that $\dim(H \cap S) = 1$. Let $\mathfrak{p}$ be the ideal of H in $\mathcal{R}(X)$ and pick any $x_0 \in \operatorname{Reg}(H)$. Then $\mathcal{R}(X)_{x_0}$ (the localization of $\mathcal{R}(X)$ at the maximal ideal $\mathfrak{m}$ of x_0, so $\mathfrak{p} \subset \mathfrak{m}$) is a factorial ring and $\operatorname{ht}(\mathfrak{p}\mathcal{R}(X)_{x_0}) = 1$, hence $\mathfrak{p}\mathcal{R}(X)_{x_0} = h\mathcal{R}(X)_{x_0}$ for some $h \in \mathfrak{p}$. Take $g_1, \ldots, g_r \in \mathcal{R}(X)$ such that $\mathfrak{p}\mathcal{R}(X) = (g_1, \ldots, g_r)$, then there exist $\lambda_i, s_i \in \mathcal{R}(X)$ with $s_i(x_0) \neq 0$ (in particular, $s_i \notin \mathfrak{p}$) such that $s_i g_i = \lambda_i h$, for $i = 1, \ldots, r$.

Consider $U = X \setminus \{s_1 \cdots s_r = 0\}$ which is Zariski open in X. Then we have $\mathfrak{p}\mathcal{R}(X)_x = (h)\mathcal{R}(X)_x$ for all $x \in U_1 = U \cap \operatorname{Reg}(H)$.

Take now $x_0 \in U_1$, we have $f_j = \rho_j h^{\alpha_j}$ with $\alpha_j \geq 0$ and $\rho_j \in \mathcal{R}(X)_{x_0}$ such that h does not divide ρ_j (in particular, $\rho_j \notin \mathfrak{p}\mathcal{R}(X)_{x_0}$). Then $\rho_j = p_j/q_j$, $j = 1, \ldots, s$, where $p_j, q_j \in \mathcal{R}(X)$ and $p_j(x_0) \neq 0, q_j(x_0) \neq 0$. Let U_2 be the Zariski open set $U_1 \setminus \{q_1 \cdots q_s = 0\} \subset H$, then for all $x \in U_2$, $q_j f_j = p_j h^{\alpha_j}$, p_j, q_j do not change sign in a neighbourhood of x, for $j = 1, \ldots, s$ and h changes sign in a neighbourhood of x, since locally h is a parameter of H in x. Hence for any $j = 1, ..., s$, f_j changes sign in a neighbourhood of x if α_j is odd and does not change sign if α_j is even.

Using the fact that $\dim(S \cap H) = 1$, $\dim(C) \leq 1$ and $s_i, p_j, q_j \notin \mathfrak{p}$, there exist a Zariski dense open set Ω in U_2 such that f_j does not change sign through Ω, for all $j = 1, \ldots, s$, then α_j is even for all $j = 1, \ldots, s$. But also there is a Zariki dense open set Ω' in U_2 such that $\Omega' \subset \overline{S} \setminus S$, then there is $l \in \{1, \ldots, s\}$ such that f_l changes sign through Ω', and α_l is odd, which is impossible. □

Proposition 1.4 *Let S be a semialgebraic set in X.*

(1) Let S be open. Then S is generically basic if and only if there exist $p_1, \ldots, p_l \in \partial_z S$ such that $S \setminus \{p_1, \ldots, p_l\}$ is basic open.

(2) S is basic open if and only if S is generically basic and $S \cap \partial_z S = \emptyset$.

(3) Let S be closed. Then S is generically basic if and only if S is basic closed.

Proof: First we prove *(1)*. The "if part" is trivial. Suppose S to be open and generically basic, that is there are $f_1, ..., f_r \in \mathcal{R}(X)$ and an algebraic set $C \subset X$ with $\dim(C) \leq 1$ such that

$$S \setminus C = \{f_1 > 0, \ldots, f_r > 0\}.$$

We suppose first that C is a curve and we can also suppose that $\{f_1 \cdots f_r = 0\} \subset C$.

$C \cap S$ is an open semialgebraic set in C, then using [Rz, 2.2], there exist $g_1 \in \mathcal{R}(C)$ such that

$$C \cap S = \{x \in C : g_1(x) > 0\}$$

$$\overline{C \cap S} = \{x \in C : g_1(x) \geq 0\}$$

choose g_1 to be the restriction of a regular function $g \in \mathcal{R}(X)$.

For each $i = 1, \dots, r$ consider the open sets $B_i = \{x \in X : f_i(x) < 0\}$ and the closed sets $T_i = (\overline{S} \cap \{g \le 0\}) \cup (\overline{B_i} \cap \{g \ge 0\})$.

Applying [BCR, 7.7.10] to T_i, f_i and g for $i = 1, ..., r$, we can find $p_i, q_i \in \mathcal{R}(X)$, with $p_i > 0$, $q_i \ge 0$, such that

(i) $F_i = p_i f_i + q_i g$ has the same signs as f_i on T_i;

(ii) The zero set $Z(q_i)$ of q_i verifies, $Z(q_i) = \mathrm{Adh_z}(Z(f_i) \cap T_i)$.

We remark the following:

a) $F_i(S \setminus C) > 0$ for $i = 1, \dots, r$, since F_i has the same signs as f_i on $T_i \cap S$ and outside is the sum of a strictly positive function and a nonnegative one.

b) $F_i(B_i) < 0$ for $i = 1, \dots, r$, by the same reasons.

c) $Z(q_i) \subset \partial_{\mathrm{z}} S$. In fact, denote

$$Z_1^i = Z(f_i) \cap \overline{S} \cap \{g \le 0\}$$

$$Z_2^i = Z(f_i) \cap \overline{B_i} \cap \{g \ge 0\}$$

then we have $Z(q_i) = \mathrm{Adh_z}(Z_1^i) \cup \mathrm{Adh_z}(Z_2^i) \subset Z(f_i) \subset C$. Since g is positive on $C \cap S$ and $Z_1^i \subset C \cap \{g \le 0\}$, we have $Z_1^i \cap S = \emptyset$, hence $Z_1^i \subset \partial(S)$. Indeed, since $B_i \cap S = \emptyset$ and S is open we have $\overline{B_i} \cap S = \emptyset$. Moreover $C \cap \overline{B_i} \subset \{g \le 0\}$, hence $Z_2^i \subset \{g = 0\} \cap C \subset \overline{S}$, then $Z_2^i \subset \partial(S)$.

From these remarks, denoting $Z = \bigcup_{i=1}^r Z(q_i)$, we have

$$S \setminus Z = \{F_1 > 0, \dots, F_r > 0\}.$$

In fact, if $x \in S \setminus C$ then $F_i(x) > 0$ for $i = 1, ..., r$, if $x \in (C \cap S) \setminus Z$ then $f_i(x) \ge 0$, $q_i(x) \ge 0$ and $g(x) > 0$, hence $F_i(x) > 0$ for $i = 1, ..., r$ and $x \in S \setminus Z$. Otherwise, suppose $x \notin S \setminus Z$, then $x \in (X \setminus S) \cup (S \cap Z)$: if $x \in X \setminus S$ there is $l \in \{1, \dots, r\}$ such that $f_l(x) \le 0$, and we can have $x \notin C$ or $x \in C$, in the first case $f_i(x) \ne 0$ for all i, so $x \in B_l$ and $F_l(x) < 0$, in the second case $g(x) \le 0$, $q_l(x) \ge 0$, so $F_l(x) \le 0$; if $x \in S \cap Z$ there is $l \in \{1, \dots, r\}$ such that $q_l(x) = 0$, then $f_l(x) = 0$ and $F_l(x) = 0$. In any way, there is l such that $F_l(X) \le 0$ if $x \notin S \setminus Z$.

By 1.3 and remark *c)* above we have that there exist $p_1, \dots, p_l \in \partial_{\mathrm{z}} S$ such that $\bigcup_{i=1}^r Z(q_i) \cap S = \{p_1, \dots, p_l\}$, hence

$$S \setminus \{p_1, \dots, p_l\} = \{F_1 > 0, \dots, F_r > 0\}.$$

If $C = \{a_1, \dots, a_m\}$ is a finite set we have to check that we can throw out from C all the a_i which do not lie in $\partial_{\mathrm{z}} S$. This can be done as before by taking the function 1 at the place of g and putting $T_i = \overline{B_i}$.

From *(1)* we have immediately *(2)*, because if S is generically basic and $\partial_{\mathrm{z}} S \cap S = \emptyset$, S is basic open, since following the proof above we have $Z(q_i) \cap S = \emptyset$ for $i = 1, ..., r$. On the countrary, if S is basic open then it is generically basic and $\partial_{\mathrm{z}} S \cap S = \emptyset$ (because $\partial_{\mathrm{z}} S \subset \{f_1 \cdots f_r = 0\}$ if $S = \{f_1 > 0, \dots, f_r > 0\}$).

Finally we prove *(3)*. The "if part" is trivial. Then suppose S to be closed and generically basic, i. e.

$$S \setminus C = \{f_1 > 0, \ldots, f_r > 0\}$$

with $f_1, \ldots, f_r \in \mathcal{R}(X)$ and $\dim(C) < 2$. We can suppose $\{f_1 \cdots f_r = 0\} \subset C$.

$C \cap S$ is a closed semialgebraic in C, by [Rz, 2.2], there is $g_1 \in \mathcal{R}(C)$ such that

$$C \cap S = \{x \in C : g_1(x) \geq 0\}.$$

Take $g \in \mathcal{R}(X)$ as above, $f \in \mathcal{R}(X)$ a positive equation of C and $T = S \cap \{g \leq 0\}$. Applying again [BCR, 7.7.10] to T, f and g, we find $p, q \in \mathcal{R}(X)$ with $p > 0$, $q \geq 0$ such that

(i) $h = pf + qg$ has the same sign as f on T;

(ii) $Z(q) = \mathrm{Adh_z}(Z(f) \cap T)$.

Notice that $h(S) \geq 0$ and $Z(f) \cap T = C \cap S \cap \{g \leq 0\} \subset \{g = 0\} \cap C$, because $C \cap S \subset \{g \geq 0\}$; but $\{g = 0\} \cap C$ is a finite set contained in S, then $Z(q)$ is a finite set contained in S.

We will prove that

$$S = \{f_1 \geq 0, \ldots, f_r \geq 0, h \geq 0\}.$$

In fact, if $x \in S \setminus C$ then $f_i(x) > 0$ for all i and $h(x) > 0$, if $x \in S \cap C$ then $f_i(x) \geq 0$ for all i (because $\dim(C) < 2$) and $h(x) \geq 0$, hence $S \subset \{f_1 \geq 0, \ldots, f_r \geq 0, h \geq 0\}$. Otherwise suppose $x \notin S \supset S \setminus C$, then there is $l \in \{1, ..., r\}$ such that $f_l(x) \leq 0$, if $x \notin C$, $f_i(x) \neq 0$ for all $i = 1, ..., r$, then $f_l(x) < 0$; if $x \in C \setminus (C \cap S)$, $f_l(x) \leq 0$, $q(x) \neq 0$ and $g(x) < 0$, then $h(x) < 0$. □

Remarks 1.5 (1) Let S be a closed semialgebraic set, then S is basic closed if and only if $S \setminus \partial_z S$ is basic open.

In fact, if S is basic closed then it is generically basic, hence $S \setminus \partial_z S$ is generically basic and as $(S \setminus \partial_z S) \cap \partial_z (S \setminus \partial_z S) = \emptyset$, $S \setminus \partial_z S$ is basic; on the contrary, if $S \setminus \partial_z S$ is basic open then S is basic closed, since S is closed.

(2) Let S be a semialgebraic set, and let S^* denote the set $\mathrm{Int}(\overline{S})$. Then, S is basic open if and only if S^* is generically basic and $S \cap \partial_z S = \emptyset$.

In fact, S and S^* are generically equal.

2 Basicness and sign distributions

We recall some definitions and results from [ABF]. Let X be a compact, non singular, real algebraic surface and $Y \subset X$ an algebraic curve.

Consider a *partial sign distribution* σ on $X \setminus Y$, which gives the sign 1 to some connected components of $X \setminus Y$ (whose union is denoted by $\sigma^{-1}(1)$) and the sign -1 to some others (whose union is denoted by $\sigma^{-1}(-1)$). So $\sigma^{-1}(1)$ and $\sigma^{-1}(-1)$ are disjoint open semialgebraic sets in X.

Definitions 2.1 (1) *A sign distribution* σ *is* completable *if* $\sigma^{-1}(1)$ *and* $\sigma^{-1}(-1)$ *can be separated by a regular function, i.e. there is* $f \in \mathcal{R}(X)$ *such that* $f(\sigma^{-1}(1)) > 0$, $f(\sigma^{-1}(-1)) < 0$ *and* $f^{-1}(0) \supset Y$. *Briefly, we say that* f induces σ.

(2) *A sign distribution* σ *is* locally completable *at a point* $p \in Y$ *if there is* $f \in \mathcal{R}(X)$ *such that* f *induces* σ *on a neighbourhood of* p.

(3) *An irreducible component* Z *of* Y *is a* type changing component *with respect to* σ *if there exist two* *nonempty open sets* $\Omega_1, \Omega_2 \subset Z \cap \mathrm{Reg}(Y)$ *such that*
(a) $\Omega_1 \subset \overline{\sigma^{-1}(1)} \cap \overline{\sigma^{-1}(-1)}$,
(b_+) $\Omega_2 \subset \mathrm{Int}(\overline{\sigma^{-1}(1)})$ *or* *(b_-)* $\Omega_2 \subset \mathrm{Int}(\overline{\sigma^{-1}(-1)})$.
If (b_+) (resp. (b_-)) holds we say that Z *is* positive type changing *(resp.* negative type changing*) with respect to* σ.

(4) *An irreducible component* Z *of* Y *is a* change component *if there exist a nonempty open set* $\Omega \subset Z \cap \mathrm{Reg}(Y)$ *verifying (a).*

Completable sign distributions are characterized by the following theorem.

Theorem 2.2 (See [ABF, 1.4 and 1.7]) *Denote by* Y^c *the union of the change components of* Y *with respect to* σ. *Then* σ *is completable if and only if*
(1) Y *does not have type changing component with respect to* σ;
(2) σ *is locally completable at any point* $p \in \mathrm{Sing}(Y)$;
(3) *There exist an algebraic curve* $Z \subset X$ *such that* $Z \cap (\sigma^{-1}(1) \cup \sigma^{-1}(-1)) = \emptyset$ *and* $[Z] = [Y^c]$ *in* $\mathrm{H}_1(X, \mathbb{Z}_2)$.
Moreover condition (2) *becomes condition* (1) *after the blowings-up of the canonical desingularization of* Y, *namely each point where* σ *is not locally completable corresponds to at least one type changing component of the exceptional divisor with respect to the lifted sign distribution.*

Proposition 2.3 *(*See [ABF, "the procedure"]) *Condition* (2) *of theorem 2.2 can be tested, without performing the blowings-up, by an algorithm that only uses the Puiseux expansions of the branches of* Y *at its singular points.*

In fact, there are two algorithms:

$\boxed{\text{A1}}$ (see [ABF, 2.4.19]) Given a branch C of an algebraic curve through a point p_0 and an integer $\rho > 0$, it is possible to find explicitely, in terms of the Puiseux expansion of C, the irreducible Puiseux parametrizations of all analytic arcs γ through p_0 with the following properties:

a) Denoting respectively by γ_ρ and by C_ρ the strict transform of γ and C after ρ blowings-up in the standard resolution of C and by D_ρ the exceptional divisor arising at the last blowing-up, then $\gamma_{\rho-1}$ is parametrized by

$$\begin{cases} x = t \\ y = at + \cdots \end{cases}$$

and $C_{\rho-1} \cap \gamma_{\rho-1} = 0 \in D_{\rho-1}$.

b) $\gamma_{\rho-1}$ and $C_{\rho-1}$ have distinct tangents at 0.

$\boxed{\text{A2}}$ (see [ABF, solution to problem 2]) Given an analytic arc γ and a region of $\mathbb{R}^2$ bounded by two analytic arcs γ_1, γ_2, with $\gamma, \gamma_1, \gamma_2$ through $0 \in \mathbb{R}^2$, it is possible to decide, looking at the Puiseux parametrizations, whether γ crosses the region or not.

Using $\boxed{\text{A1}}$ and $\boxed{\text{A2}}$ one can decide whether D_ρ is positive or negative type changing with respect to the lifted sign distribution σ_ρ without performing the blowings up, because for the family of arcs given by $\boxed{\text{A1}}$, whose strict transform are transversal to D_ρ, we can decide by $\boxed{\text{A2}}$ whether or not σ changes sign along some elements of the family and whether or not σ has constant positive or constant negative sign along some other ones.

Now let S be an open semialgebraic set.

Notation 2.4 Let $A_1, \ldots, A_t$ be the connected component of $X \setminus (S \cup \partial_{\mathrm{z}} S)$. For each $i = 1, \ldots, t$ we denote by σ_i^S (or simply σ_i when there is not risk of confusion) the following sign distribution on $X \setminus \partial_{\mathrm{z}} S$:

$$\begin{aligned} (\sigma_i^S)^{-1}(1) &= S \setminus \partial_{\mathrm{z}} S \\ (\sigma_i^S)^{-1}(-1) &= A_i \end{aligned}$$

Lemma 2.5 *Let S be a semialgebraic set and $S^* = \mathrm{Int}(\overline{S})$. Then,* $\dim(S^* \cap \partial_{\mathrm{z}} S^*) = 1$ *if and only if there exists $i \in \{1, \ldots, t\}$ such that $\partial_{\mathrm{z}} S$ has a positive type changing component with respect to σ_i^S.*

Proof: Suppose $\dim(S^* \cap \partial_{\mathrm{z}} S^*) = 1$, then there is an irreducible component H of $\partial_{\mathrm{z}} S^*$ such that $\dim(H \cap S^*) = 1$. So we can find an open 1-dimensional set $\Omega \subset H \cap S^*$, hence

$$\Omega \subset \mathrm{Int}(\overline{S}) = \mathrm{Int}(\overline{S \setminus \partial_{\mathrm{z}} S}) = \mathrm{Int}(\overline{\sigma_i^{-1}(1)})$$

for each $i = 1, \ldots, t$. And we can find another open 1-dimensional set $\Omega' \subset \partial(S^*) \cap H$ such that

$$\Omega' \subset \overline{S^*} = \overline{S} = \overline{S \setminus \partial_{\mathrm{z}} S} = \overline{\sigma_i^{-1}(1)}$$

for each $i = 1, \ldots, t$ and

$$\Omega' \subset X \setminus S^* = \bigcup_{i=1}^{t} \overline{A_i} \; .$$

Then $\Omega' \subset \bigcup_{i=1}^{t}(\overline{A_i} \setminus A_i)$, since $A_i \cap S = \emptyset$ and S is open. Hence, there exist a 1-dimensional open set $\Omega'' \subset \Omega'$ and $i_0 \in \{1, ..., t\}$ such that $\Omega'' \subset \overline{A_{i_0}} \setminus A_{i_0}$, because $\dim(\Omega') = 1$ and we have a finite number of A_i. Hence $\Omega'' \subset \overline{\sigma_{i_0}^{-1}(1)} \cap \overline{\sigma_{i_0}^{-1}(-1)}$. But since H is a 1-dimensional component of $\partial_{\mathrm{z}} S^*$ and $\partial_{\mathrm{z}} S^* \subset \partial_{\mathrm{z}} S$, H is an irreducible component of $\partial_{\mathrm{z}} S$ of dimension 1; then if we take $\Omega_1 = \Omega'' \cap \mathrm{Reg}(\partial_{\mathrm{z}} S)$ and $\Omega_2 = \Omega \cap \mathrm{Reg}(\partial_{\mathrm{z}} S)$, H is a positive type changing component with respect to σ_{i_o}.

On the contrary suppose that H is an irreducible component of $\partial_{\mathrm{z}} S$ which is a positive type changing component with respect to some σ_l $(l = 1, ..., t)$. So there exist open sets $\Omega_1, \Omega_2 \subset H$ of $\mathrm{Reg}(\partial_{\mathrm{z}} S)$ such that $\Omega_1 \subset \overline{\sigma_l^{-1}(1)} \cap \overline{\sigma_l^{-1}(-1)}$ and $\Omega_2 \subset \mathrm{Int}(\overline{\sigma_l^{-1}(1)})$. Then $\Omega_2 \subset S^*$ and $\dim(S^* \cap H) = 1$. But $\Omega_1 \subset \overline{S \setminus \partial_{\mathrm{z}} S} = \overline{S^*}$ and $\Omega_1 \subset \overline{A_i}$, then $\Omega_1 \subset \overline{S^*} \setminus S^*$ because $X \setminus S^* = \bigcup \overline{A_i}$. So H is an irreducible component of $\partial_{\mathrm{z}} S^*$, hence $\dim(S^* \cap \partial_{\mathrm{z}} S^*) = 1$.

□

Proposition 2.6 *Let S be a semialgebraic set in the sphere $\mathbb{S}^2$ such that $\partial_{\mathrm{z}} S \cap S = \emptyset$ (resp. a closed semialgebraic set in $\mathbb{S}^2$). Then S is basic open (resp. closed) if and only if for each $i = 1, \ldots, t$, σ_i^S is completable.*

Proof: It suffices to prove the result for a semialgebraic S such that $\partial_{\mathrm{z}} S \cap S = \emptyset$, because if S is closed, applying 1.5 we have done.

Suppose S to be basic open, then S^* is generically basic and $\dim(S^* \cap \partial_{\mathrm{z}} S^*) < 1$. By lemma 2.5 no irreducible component of $\partial_{\mathrm{z}} S$ is positive type changing with respect to σ_i^S for each $i = 1, ..., t$. But also they cannot be negative type changing, because any curve in $\mathbb{S}^2$ divides $\mathbb{S}^2$ into connected components in such a way that none of them lies on both sides of a branch of the curve (because the curves in $\mathbb{S}^2$ have orientable neighbourdhoods).

If for some $i \in \{1, ..., t\}$, σ_i^S were not locally completable at some point $p \in \partial_{\mathrm{z}} S$, we could find (Theorem 2.2) a non singular surface V together with a contraction $\pi : V \longrightarrow \mathbb{S}^2$ of an algebraic curve $E \subset V$ to the point p, with $\pi^{-1}(\partial_{\mathrm{z}} S)$ normal crossing in V, such that an irreducible component D of E would be type changing with respect to $\sigma'_i = \sigma_i^S \cdot \pi$. But $p \notin S \cup A_i$, so σ'_i is defined on $V \setminus \pi^{-1}(\partial_{\mathrm{z}} S)$ by

$$\begin{array}{rcl} (\sigma'_i)^{-1}(1) = & \pi^{-1}(S) = & T \\ (\sigma'_i)^{-1}(-1) = & \pi^{-1}(A_i) & ; \end{array}$$

and as $\pi : V \setminus (\pi^{-1}(\partial_{\mathrm{z}} S)) \rightarrow \mathbb{S}^2 \setminus \partial_{\mathrm{z}} S$ is a biregular isomorphism, $\sigma'_i = \sigma_i^T$. So T and T^* are generically basic, being biregularly isomorphic to S, hence $\dim(\partial_{\mathrm{z}} T^* \cap T^*) < 1$; so D cannot be positive type changing by lemma 2.5. We have to exclude that D is negative type changing. Suppose it is so; then we can find two open sets $\Omega, \Omega' \subset D$ such that

$\pi^{-1}(A_i)$ lies on both sides of Ω and Ω' divides T from $\pi^{-1}(A_i)$. Then there exists an irreducible component Z of $\partial_z S$ such that its strict transform Z' crosses D between Ω and Ω'. Then Z' must cross $\pi^{-1}(A_i)$, because this open set is connected. This is impossible since A_i does not lie on both sides of Z. Then no irreducible component of $\partial_z S$ is type changing with respect to σ_i^S and σ_i^S is locally completable at any $p \in \partial_z S$ for all i; so σ_i^S is completable, since $\mathrm{H}_1(\mathbb{S}^2, \mathbb{Z}_2) = 0$ (2.2).

Suppose now that σ_i^S is completable for each $i = 1, ..., t$ and let $f_i \in \mathcal{R}(\mathbb{S}^2)$ be a regular function inducing σ_i^S. Clearly $S \subset \{f_1 > 0, \dots, f_t > 0\}$. But if $x \notin S$ then $x \in A_l$ for some $l = 1, ..., t$ or $x \in \partial_z S$. If $x \in A_l$, then $f_l < 0$; if $x \in \partial_z S$, then $f_i(x) = 0$ for all $i = 1, ..., t$, since f_i vanishes on $\partial_z S$ by the very definition of completability. So $S = \{f_1 > 0, \dots, f_t > 0\}$ and it is basic. □

Before proving an analogous result for a general surface we need a lemma.

Lemma 2.7 *Let $S \subset X$ be an open semialgebraic set such that $S = S^*$, $\partial_z S \cap S = \emptyset$ and $\partial_z S$ is normal crossing. Let H be an irreducible component of $\partial_z S$. Then there exist a non singular algebraic set $H' \subset X$ such that*

(a) $[H] = [H']$ in $\mathrm{H}_1(X, \mathbb{Z}_2)$,

(b) H and H' are transversals,

(c) $H' \cap S = \emptyset$.

Proof: H is a smooth compact algebraic curve, composed by several ovals and locally disconnects its neighbourhood; moreover S is locally only on one side of H, because $S = S^*$ and $H \cap S^* = \emptyset$, at least outside a neighbourhood of the singular points lying in H, namely the points where H crosses an other component of $\partial_z S$. At each of these points we have only two possible situations (see figures 2.7). From this it is clear how to construct a smooth differentiable curve C_H with $[C_H] = [H]$ in $\mathrm{H}_1(X, \mathbb{Z}_2)$ such that C_H is transversal to each irreducible component of $\partial_z S$ and $C_H \cap S = \emptyset$, using a suitable tubular neighbourhood of H.



Figure 2.7.a



Figure 2.7.b

Now we want to approximate C_H by a nonsingular algebraic curve H' with the same properties. To do this we use the fact that $[H]$ gives a strongly algebraic line bundle

$\pi : E \to X$ on X (see [BCR, 12.2.5]). For this line bundle, H is the zero set of an algebraic section h and C_H is the zero set of a $\mathcal{C}^\infty$ section c. Let $Q_1, \dots, Q_k$ be the points in the set $H \cap C_H$. We can take a finite open covering $V_1, \dots, V_l$ of X with the following properties:

1) $Q_i \in V_i$ for $i = 1, ..., k$ and $V_1, \dots, V_k$ are pairwise disjoint.

2) For each $j = 1, ..., l$ there exist an algebraic section s_j of E such that $s_j(x) \neq 0$ for each $x \in V_j$ (s_j generates E_x for each $x \in V_j$).

Take a $\mathcal{C}^\infty$ partition of the unity $\{\varphi_1, \dots, \varphi_l\}$ associated to the covering, with the property that for $i = 1, ..., k$, $\varphi_i^{-1}(1)$ is a closed neighbourhood of Q_i in V_i, $\varphi_i(Q_j) = 0$ for $j \neq i$. For $j = 1, ..., l$ we can write $c|_{V_j} = \alpha_j s_j$, with $\alpha_j \in \mathcal{C}^\infty(V_j)$ and $\alpha_j(Q_j) = 0$ if $j = 1, ..., k$, because s_j generates the fiber. Then we have

$$c = \sum_{j=1}^{l} \varphi_j c = \sum_{j=1}^{l} (\varphi_j \alpha_j) s_j$$

and the smooth functions $\beta_j = \varphi_j \alpha_j \in \mathcal{C}^\infty(X)$ vanishes at Q_i for $i = 1, ..., k$.

By a classical relative approximation theorem (see [BCR, 12.5.5]) we can approximate β_j on X by a regular function f_j on X vanishing at $\{Q_1, \dots, Q_k\}$. Then the algebraic section

$$s = \sum_{j=1}^{l} f_j s_j$$

has an algebraic zero set H' passing through $Q_1, \dots, Q_k$. Moreover $H' \cap S = \emptyset$, because H' is very close to C_H, and $[H'] = [H]$ in $\mathrm{H}_1(X, \mathbb{Z}_2)$. □

Finally we have.

Theorem 2.8 *Let X be a compact, non-singular, real algebraic surface and $S \subset X$ be a semialgebraic set with $\partial_z S \cap S = \emptyset$ (resp. $S \subset X$ be a closed semialgebraic set).*

Then S is basic if and only if for each $i = 1, \dots, t$ the sign distribution σ_i^S verifies the following two properties:

(a) No irreducible component of $\partial_z S$ is positive type changing with respect to σ_i^S.

(b) No irreducible component of the exceptional divisor of a standard resolution $\pi : V \to X$ of $\partial_z S$ is positive type changing with respect to $\sigma_i' = \sigma_i^S \cdot \pi$.

Proof: It suffices to prove for S with $S \cap \partial_z S = \emptyset$, because this implies that S is open and by remark 1.5.1 we have done for S closed.

The "only if part" is the same as for $\mathbb{S}^2$, without the argument proving there are not negative type changing component. For the "if part" we can reason as follows.

Let $\pi : V \to X$ be the standard resolution of $\partial_z S$. Denote by Y the curve $\pi^{-1}(\partial_z S)$ (see [EC] or [BK]), Y is normal crossing in V. Each irreducible component of Y is a non-singular curve consisting possibly of several ovals. Each of this ovals can have an orientable neighbourhood (in which case it is homologically trivial) or a non orientable neighbourhood isomorphic to the Möbius band.

Denote as it is usual σ'_i the sign distribution $\sigma_i^S \cdot \pi$ and consider the sets $T = \pi^{-1}(S)$ and $T^* = \mathrm{Int}(\overline{T})$. Conditions *(a)* and *(b)* for σ_i^S imply that Y has not positive type changing components with respect to σ'_i. So by 2.5, $\partial_{\mathrm{z}} T^* \cap T^*$ is a finite set, and as $\partial_{\mathrm{z}} T^* \subset Y$ has not isolated points and T^* is open, we have $T^* \cap \partial_{\mathrm{z}} T^* = \emptyset$.

Consider now the sign distributions $\sigma_j^{T^*}$ on $V \setminus \partial_{\mathrm{z}} T^*$ defined as before for $j = 1, \dots, l$, where l is the number of connected component of $V \setminus (T^* \cup \partial_{\mathrm{z}} T^*)$; clearly they do not have positive type changing components. Apply 2.7 to each irreducible component H of $\partial_{\mathrm{z}} T^*$ being a change component for some $\sigma_j^{T^*}$. Then, we find a non-singular algebraic set H' such that $H' \cap T^* = \emptyset$ and $[H] = [H']$ in $\mathrm{H}_1(V, \mathbb{Z}_2)$. The union of all H' gives an algebraic set $Z \subset V$.

Remark that if H is a negative type changing component with respect some $\sigma_j^{T^*}$ then this phenomenon occurs along an oval of H whose neighbourhood is a Möbius band, because in the other case the two sides of the oval whould be in different connected components of $V \setminus \partial_{\mathrm{z}} T^*$. Take the sign distributions τ_K on $V \setminus (\partial_{\mathrm{z}} T^* \cup Z)$, $k = 1, \dots, m$, defined by

$$\begin{aligned} (\tau_k)^{-1}(1) &= T^* \\ (\tau_k)^{-1}(-1) &= B_k, \end{aligned}$$

where $B_1, \dots, B_m$ are the connected components of $V \setminus (T^* \cup \partial_{\mathrm{z}} T^* \cup Z)$.

We claim that τ_k is completable for each $k = 1, ..., m$. In fact, we prove that conditions (1), (2) and (3) of 2.2 are verified.

(1) No irreducible component of $\partial_{\mathrm{z}} T^* \cup Z$ is neither positive nor negative type changing with respect τ_k: this is true because now the sign -1 can occur at most on one side of each irreducible component H of $\partial_{\mathrm{z}} T^* \cup Z$ (a Möbius band is divided by two transversal generators of its not vanishing homological class into two connected components), then there are not negative type changing components with respect τ_k.

(2) τ_k is completable at each point $p \in \partial_{\mathrm{z}} T^* \cup Z$. In fact, if $\partial_{\mathrm{z}} T^* \cup Z$ is normal crossing in p, since there are not type changing components, τ_k is locally completable at p. But, in general $\partial_{\mathrm{z}} T^* \cup Z$ is not normal crossing. If p_0 is not normal crossing we have, by construction 2.7, two irreducible components H, H_1 of $\partial_{\mathrm{z}} T^*$ and one irreducible component H' of Z, with $[H'] = [H]$, meeting pairwise transversally at p_0.



Figure 2.8

Then by construction we have two signs $+1$ between H and H_1 near p_0 (if not H' would not cross H) and at most two signs -1 between H and H' or between H' and H_1 (see figure 2.8). So τ_k is locally completable at p_0.

(3) If H is a change component for τ_k, then $H \subset \partial_z T^*$ and if $[H] \neq 0$ then for some irreducible component Z_H of Z, $[H \cup Z_H] = 0$ and $Z_H \cap T^* = \emptyset$, $Z_H \cap B_k = \emptyset$, by construction.

So τ_k is completable for $k = 1, ..., m$. Let P_k be a regular function inducing τ_k. Then

$$T^* \subset \{P_1 > 0, \ldots, P_m > 0\},$$

but if $x \notin T^*$, $x \in (\bigcup_{k=1}^m B_k) \cup \partial_z T^* \cup Z$, hence at least one among P_k verifies $P_k(x) \leq 0$. So

$$T^* = \{P_1 > 0, \ldots, P_m > 0\}$$

then it is basic, hence S is generically basic, but $S \cap \partial_z S = \emptyset$ so, by 1.4, S is basic. □

From 2.8 we find for surfaces a geometric proof of a general basicness characterization in [AR2]. We call *birational model* of a semialgebraic set S any semialgebraic set obtained from S by a birational morphism on X.

Corollary 2.9 *Let X be a surface and $S \subset X$ a semialgebraic set. Then, S is basic open if and only if $\partial_z S \cap S = \emptyset$ and for each birational model T of S we have $\partial_z T^* \cap T^*$ is a finite set.*

Proof: By 1.4 we are done the *only if part.* Suppose now that $\partial_z S \cap S = \emptyset$ and for each birational model T of S we have that $\partial_z T^* \cap T^*$ is a finite set. Then, S is open and we will prove that it is basic open.

Take a compactification of X (for instance its closure in a projective space) and then take a non-singular birational model X_1 of X, obtained by a finite sequence of blowings-up along smooth centers. The strict transform S_1 of S is a birational model of S.

Consider now the standard resolution $\pi : X_2 \to X_1$ of $\partial_z S_1$ and take S_2 the strict transform of S_1 by π. Then $\partial_z S_2$ is normal crossing and $\partial_z S_2^* \cap S_2^*$ is a finite set, because S_2 is a birational model of S. But, $\partial_z S_2^* \subset \partial_z S_2$ has not isolated points (it is normal crossing) and S_2^* is open, then $\partial_z S_2^* \cap S_2^* = \emptyset$. So by 2.5 and 2.8, S_2^* is basic open. Hence S is generically basic and, as $\partial_z S \cap S = \emptyset$, S is basic open. □

3 Geometric review of fans

For all notions of real algebra, real spectra, specialization, real valuation rings, etc., we refer to [BCR]. Only for tilde operation we use a slightly different definition: for a semialgebraic set S in an algebraic set X, $\tilde{S}$ is the constructible set of $\mathrm{Spec}_r(\mathcal{R}(X))$ (instead of $\mathcal{P}(X)$) defined by the same formula which defines S. The properties of this tilde operation are the the same as the usual ones (see [BCR, chap.7]).

Let K be a real field, a subset $F = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ of $\mathrm{Spec}_r K$ is a *4-element fan* (or simply a *fan*) if each α_i is the product of the other three, that is for each $f \in K$ we have

$$\alpha_i \alpha_j \alpha_k(f) = \alpha_l(f)$$

for all $\{i,j,k,l\} = \{1,2,3,4\}$, where $\alpha(f)$ denotes the sign (1 or -1) of f in the ordering $\alpha \in \mathrm{Spec}_r(K)$.

Given a fan F we can find a valuation ring V of K such that

a) Each $\alpha_i \in F$ is compatible with V; that is, the maximal ideal $\mathfrak{m}_V$ of V is α_i-convex.

b) F induces at most two orderings in the residue field k_V of V.

In this situation we say that F trivializes along V (see [BCR, chap.10] and [Br]).

Let X be a real algebraic set, and $\mathcal{K}(X)$ be the function field of X, that is a finitely generated real extension of $\mathbb{R}$. Denote $K = \mathcal{K}(X)$.

Definition 3.1 *A fan F of K is* associated to a real prime divisor V *if*

(a) *V is a discrete valuation ring such that F trivializes along V.*

(b) *The residue field k_V of V is a finitely generated real extension of $\mathbb{R}$ such that* $\mathrm{dg.tr.}[K:\mathbb{R}] = \mathrm{dg.tr.}[k_V:\mathbb{R}] + 1$.

Remark 3.2 Let F be a not trivial fan (i.e. the α_i's are distincts) associated to a real prime divisor V, then it induces two distinct orderings τ_1, τ_2 in k_V ([BCR, 10.1.10]). If $F = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ we suppose that α_1, α_3 (resp. α_2, α_4) induce τ_1 (resp. τ_2) and we write this

$$\begin{tikzcd}
V \arrow[d] & \alpha_1 \arrow[dr] & & \alpha_3 \arrow[dl] & \alpha_2 \arrow[dr] & & \alpha_4 \arrow[dl] \\
k_V & & \tau_1 & & & \tau_2 &
\end{tikzcd}$$

Conversely, let $\tau_1, \tau_2 \in \mathrm{Spec}_r(k_V)$ be distinct, and let $t \in V$ be a uniformizer for V. Each $f \in V$ can be written as $f = t^n u$, where n is the valuation of f and u is a unit in V. Denote by $\overline{u}$ the class of u in k_V and consider the orderings in K defined as follows:

$$\begin{array}{lcl}
\alpha_1(f) = \tau_1(\overline{u}) & ; & \alpha_3(f) = (-1)^n \tau_1(\overline{u}) \\
\alpha_2(f) = \tau_2(\overline{u}) & ; & \alpha_4(f) = (-1)^n \tau_2(\overline{u})
\end{array}$$

They form a fan F of K associated to the real prime divisor V.

We may consider $\tau_1, \tau_2 \in \mathrm{Spec}_r(k_V)$ as elements of $\mathrm{Spec}_r(V)$ with $\mathfrak{m}_V$ as support.

Then we have that α_1, α_3 (resp. α_2, α_4) specialize to τ_1 (resp. τ_2) in $\mathrm{Spec}_r(V)$.

When α specializes to τ, we write $\alpha \rightarrow \tau$.

From now we consider the field of rational functions $\mathcal{K}(X)$ of a compact non-singular real algebraic surface X, which is a finitely generated real extension of $\mathbb{R}$ with transcendence degree over $\mathbb{R}$ equal to 2.

Remark 3.3 Let F be a fan in $\mathcal{K}(X)$ associated to a real prime divisor V of $\mathcal{K}(X)$. Then, $\mathcal{R}(X) \subset V$ (because X is compact); consider the real prime ideal $\mathfrak{p} = \mathcal{R}(X) \cap \mathfrak{m}_V$. We have that V dominates $\mathcal{R}(X)_{\mathfrak{p}}$ and there are two possibilities:

1) If the height of $\mathfrak{p}$ is 1, it is the ideal of an irreducible algebraic curve $H \subset X$. Since X is non-singular, $\mathcal{R}(X)_{\mathfrak{p}}$ is a discrete valuation ring, which is dominated by V. Hence $V = \mathcal{R}(X)_{\mathfrak{p}}$ and k_V is the function field $\mathcal{K}(H)$ of H; so $\tau_1, \tau_2 \in \mathrm{Spec}_r(H)$.

2) If $\mathfrak{p}$ is a maximal ideal, it is the ideal of a point $p \in X$, because X is compact.

Definition 3.4 *Let F be a fan of $\mathcal{K}(X)$ associated to a real prime divisor V and let $\mathfrak{p} = \mathcal{R}(X) \cap \mathfrak{m}_V$. The* center of F *is the zero set $Z(\mathfrak{p})$ of $\mathfrak{p}$.*

We say that F is centered at a curve (resp. a point) *if $\mathfrak{p}$ has height 1 (resp. is maximal).*

Lemma 3.5 *Let $S \subset X$ be an open semialgebraic set. Then the following facts are equivalent:*

(i) *For each fan F of $\mathcal{K}(X)$ centered at a curve, $\#(F \cap \tilde{S}) \neq 3$*

(ii) *$\partial_z S$ has not positive type changing components with respect to the sign distributions σ_i^S for $i = 1, ..., t$, defined in 2.4.*

Proof: Suppose to have a fan F centered at a curve $H \subset X$, such that $\#(F \cap \tilde{S}) = 3$. Then by remarks 3.2 and 3.3 we have:

a) F is associated to a real prime divisor $V = \mathcal{R}(X)_{\mathfrak{p}}$, where $\mathfrak{p}$ is the ideal of H.

b) If $F = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$, then $\alpha_1, \alpha_3 \rightarrow \tau_1$, $\alpha_2, \alpha_4 \rightarrow \tau_2$ in $\mathrm{Spec}_r(V)$, with $\tau_1 \neq \tau_2$ and $\tau_1, \tau_2 \in \mathrm{Spec}_r(\mathcal{K}(H))$.

Suppose $\alpha_1, \alpha_2, \alpha_3 \in \tilde{S}$ and $\alpha_4 \notin \tilde{S}$. Remark that an element of $\mathrm{Spec}_r(\mathcal{R}(X)_{\mathfrak{p}})$ is a prime cone of $\mathrm{Spec}_r(\mathcal{R}(X))$ which support is contained in $\mathfrak{p}$. So we can consider $\alpha_i, \tau_j \in \mathrm{Spec}_r(\mathcal{R}(X))$ $(i = 1, 2, 3, 4, j = 1, 2)$ with $\tau_1, \tau_2 \in \tilde{H}$ and $\alpha_1, \alpha_3 \rightarrow \tau_1$, $\alpha_2, \alpha_4 \rightarrow \tau_2$ in $\mathrm{Spec}_r(\mathcal{R}(X))$.

We have $\tau_1 \in \overline{\tilde{S}} = \tilde{\overline{S}}$, because $\alpha_1, \alpha_3 \in \tilde{S}$. But by [BCR, 10.2.8] there are precisely two prime cone different from τ_1 in $\mathrm{Spec}_r(\mathcal{R}(X))$ specializing to τ_1, so they are α_1, α_3. And as $\alpha_1, \alpha_3, \tau_1 \in \tilde{\overline{S}}$, we get that τ_1 is an interior point of $\tilde{\overline{S}}$, so $\tau_1 \in \tilde{S}^*$. This means $\tau_1 \in \tilde{H} \cap \tilde{S}^*$, so $\dim(H \cap S^*) = 1$.

Now $\tau_2 \in \tilde{\overline{S}}$, because $\alpha_2 \in \tilde{S}$ and $\alpha_2 \to \tau_2$. Again [BCR, 10.2.8] the prime cones specializing to τ_2 and different from it are precisely α_2, α_4. Since $\alpha_4 \notin \tilde{S}$ and $\tilde{S} \cap \mathrm{Spec}_r(\mathcal{K}(X)) = \tilde{\overline{S}} \cap \mathrm{Spec}_r(\mathcal{K}(X))$, we have that $\alpha_4 \notin \tilde{\overline{S}}$, so τ_2 is not interior to $\tilde{\overline{S}}$, that is $\tau_2 \notin \tilde{S}^*$. But $\overline{S} = \overline{S^*}$, then

$$\tau_2 \in \widetilde{\overline{S^*} \setminus S^*} \subset \widetilde{\partial_{\mathrm{z}} S^*} \,.$$

This implies that $\dim(H \cap \partial_{\mathrm{z}} S^*) = 1$, then H is an irreducible component of $\partial_{\mathrm{z}} S^*$. So $\dim(S^* \cap \partial_{\mathrm{z}} S^*) = 1$ and by 2.5 $\partial_{\mathrm{z}} S$ has a positive type changing component with respect σ_i^S for some $i = 1, ..., t$.

Conversely, let H be a irreducible component of $\partial_{\mathrm{z}} S$ which is positive type changing with respect σ_i^S for some i. Then we can find open sets $\Omega_1, \Omega_2 \in H \cap \mathrm{Reg}(\partial_{\mathrm{z}} S)$ such that

a) $\Omega_1 \subset \overline{\sigma_i^{-1}(1)} \cap \overline{\sigma_i^{-1}(-1)} = \overline{S} \cap \overline{A}_i$

b) $\Omega_2 \subset \mathrm{Int}(\overline{\sigma_i^{-1}(1)}) = S^*$

Let $\mathfrak{p}$ be the ideal of H in $\mathcal{R}(X)$ and V be the discrete valuation ring $\mathcal{R}(X)_{\mathfrak{p}}$. Consider two orderings $\tau_1, \tau_2 \in \mathrm{Spec}_r(\mathcal{K}(H))$, with $\tau_1 \in \tilde{\Omega}_1$, $\tau_2 \in \tilde{\Omega}_2$, and let F be the fan defined by τ_1, τ_2 as in 3.2. So $\alpha_1, \alpha_3 \to \tau_1$, $\alpha_2, \alpha_4 \to \tau_2$ in $\mathrm{Spec}_r(\mathcal{R}(X))$, as before.

We have $\alpha_2, \alpha_4 \in \tilde{S}^*$, because $\tau_2 \in \tilde{S}^*$ and S^* is open. But $\alpha_2, \alpha_4 \in \mathrm{Spec}_r(\mathcal{K}(X))$ and $\tilde{S} \cap \mathrm{Spec}_r(\mathcal{K}(X)) = \tilde{S}^* \cap \mathrm{Spec}_r(\mathcal{K}(X))$, then $\alpha_2, \alpha_4 \in \tilde{S}$.

On the other hand, $\tau_1 \in \tilde{\overline{S}} \cap \tilde{\overline{A}}_i$. So there exist $\alpha \in \tilde{S}$ and $\beta \in \tilde{A}_i$ with $\alpha, \beta \to \tau_1$. Again by [BCR, 10.2.8] we must have $\alpha = \alpha_1$ and $\beta = \alpha_3$. So $\#(F \cap \tilde{S}) = 3$. $\square$

Lemma 3.6 *Let S be a open semialgebraic set in X such that $\partial_{\mathrm{z}} S^* \cap S^*$ is a finite set. Fix $p \in \partial S$. Then the following facts are equivalent:*

(i) *For each fan F centered at p, $\#(F \cap \tilde{S}) \neq 3$.*

(ii) *For each contraction $\pi : X' \to X$ of a curve E to the point p, no irreducible component of E is positive type changing with respect to $\sigma_i' = \sigma_i^S \cdot \pi$, for $i = 1, ..., t$.*

Proof: Suppose that there exist a contraction $\pi : X' \to X$ of a curve E to the point p and $i = 1, ..., t$ such that an irreducible component H of E is positive type changing with respect to σ_i'. But if $T = \pi^{-1}(S)$, then $(\sigma_i')^{-1}(1) = T$, $(\sigma_i')^{-1}(-1) = \pi^{-1}(A_i)$; moreover, as $p \notin S$, the set $\{\pi^{-1}(A_i) : i = 1, ..., t\}$ is precisely the set of connected component of $X' \setminus (T \cup \partial_{\mathrm{z}} T)$, then $\sigma_i' = \sigma_i^T$. Now by 3.5 there exists a fan $F = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ of $\mathcal{K}(X')$ with center the curve H such that $\#(F \cap \tilde{T}) = 3$.

The contraction π induces a field isomorphism

$$\pi_* : \mathcal{K}(X) \to \mathcal{K}(X')$$

and an injective ring homomorphism $\pi_*|_{\mathcal{R}(X)} : \mathcal{R}(X) \to \mathcal{R}(X')$. Let G be the fan of $\mathcal{K}(X')$ inverse image of F by π_*, namely

$$G = \{\pi_*^{-1}(\alpha_1), \pi_*^{-1}(\alpha_2), \pi_*^{-1}(\alpha_3), \pi_*^{-1}(\alpha_4)\} \subset \mathrm{Spec}_r(\mathcal{K}(X))$$

Then $\#(G \cap \tilde{S}) = 3$ and we have to prove that p is the center of G. Let V be the real prime divisor associated to F, then $\pi_*^{-1}(V) = W$ is the real prime divisor associated of G, so

$$\mathfrak{m}_W \cap \mathcal{R}(X) = \pi_*^{-1}(\mathcal{J}(H))$$

where $\mathcal{J}(H)$ denotes the ideal of H in $\mathcal{R}(X')$. Hence,

$$\mathfrak{m}_W \cap \mathcal{R}(X) = \mathcal{J}(\pi(H)) = \mathfrak{m}_p$$

with $\mathfrak{m}$ the maximal ideal of p.

On the contrary, we suppose that no irreducible component of E is positive type changing with respect to $\sigma_i' = \sigma_i \cdot \pi$, for each contraction π of a curve to p. Take a neighbourhood U of p, homeomorphic to a disk in $\mathbb{R}^2$ (X is non-singular), such that U does not meet any irreducible component of $\partial_z S$ unless it contains p. Consider the sign distributions δ_j in $X \setminus (\partial_z S \cup \partial U)$ for $j = 1, ..., l$, defined by

$$\begin{aligned} \delta_j^{-1}(1) &= U \cap S \\ \delta_j^{-1}(-1) &= B_j \end{aligned}$$

where $B_1, \ldots, B_l$ are the connected components of $U \setminus (S \cup \partial_z S)$. As $\partial_z S^* \cap S^*$ is a finite set, by 2.5 $\partial_z S$ has not positive type changing components with respect to σ_i^S for all $i = 1, ..., t$. We claim that $\partial_z(U \cap S)$ has no type changing components at all. In fact, as $B_j \subset A_i$ for some i, there are not positive ones; ∂U cannot be type changing because the signs may lie only on one side of it and no other component can be negative type changing, for the same reasons as in the proof of 2.6.

Now if $\partial_z S$ is normal crossing at p, by 2.2 δ_j is locally completable at p for $i = 1, ..., l$. If not, consider the standard singularity resolution $\pi : X' \to X$ of $\partial_z S$ at p and the sign distributions

$$\begin{aligned} \sigma_i' = \sigma_i^S \cdot \pi, \ \mathit{for}\ i = 1, ..., t \\ \delta_j' = \delta_j \cdot \pi, \ \mathit{for}\ i = 1, ..., l \end{aligned}$$

As no irreducible component of $\pi^{-1}(p)$ is positive type changing with respect to σ_i' for all i, the same is true with respect to δ_j' ($j = 1, ..., l$). More over each such component has a Möbius neighbourhood in $\pi^{-1}(U)$ where the sign minus can occur locally only on one side of the curve, so it cannot be negative type changing. Then, by 2.2 δ_j is locally completable at p for each $j = 1, .., l$.

Hence, for each $j = 1, ..., l$, take $f_j \in \mathcal{R}(X)$ and an open set $U_j \ni p$, $U_j \subset U$, such that f_j induces δ_j on U_j. Consider $A = \bigcap_{j=1}^{l} U_j$, then

$$S \cap A = \{f_1 > 0, \ldots, f_l > 0\} \cap A.$$

In fact, by definition of locally completable we have

$$S \cap A \subset \{f_1 > 0, \ldots, f_l > 0\};$$

but if $x \in A \setminus S$, then $x \in (\bigcup B_j) \cup \partial_z S$, so there is a j_0 such that $f_{j_0}(x) \leq 0$.

Let $F = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ a fan centered at p. Clearly $F \subset \tilde{A}$, because each α_i specializes in $\mathrm{Spec}_r(\mathcal{R}(X))$ to the prime cone having $\mathfrak{m}_p$ as support, i.e the unique prime cone giving to $f \in \mathcal{R}(X)$ the sign of $f(p)$ ([BCR, 10.2.3]). Suppose that $\alpha_1, \alpha_2, \alpha_3 \in \tilde{S}$ and $\alpha_4 \notin \tilde{S}$; then $\alpha_1, \alpha_2, \alpha_3 \in \tilde{A} \cap \tilde{S}$ and $\alpha_4 \notin \tilde{A} \cap \tilde{S}$. So for all $j = 1, ..., l$, $\alpha_i(f_j) > 0$ for $i = 1, 2, 3$ and there is j_0 such that $\alpha_4(f_{j_0}) < 0$. But this is imposible because F is a fan and $\alpha_1\alpha_2\alpha_3(f_{j_0}) \neq \alpha_4(f_{j_0})$. □

Remark 3.7 Let S be an open semialgebraic set such that $\partial_z S^* \cap S^*$ is a finite set. Let $p \in \partial_z S$ such that $p \notin \partial S$ or $\partial_z S$ is normal crossing at p. Then for each fan centered at p, $\#(F \cap \tilde{S}) \neq 3$.

Theorem 3.8 (See [Br] and [AR1]) *Let X be a real irreducible algebraic surface. Let S be a semialgebraic set such that $\partial_z S \cap S = \emptyset$ (resp. a closed semialgebraic set). Then, S is basic open (resp. basic closed) if and only if for each fan F of $\mathcal{K}(X)$ which is associated to a real prime divisor, $\#(F \cap \tilde{S}) \neq 3$.*

Proof: Suppose S to be basic open, then

$$S = \{f_1 > 0, \ldots, f_r > 0\}, \text{ with } f_1, \ldots, f_r \in \mathcal{R}(X).$$

Let $F = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ be a fan in $\mathcal{K}(X)$ and suppose that $\alpha_1, \alpha_2, \alpha_3 \in S$ and $\alpha_4 \notin S$. Then for all $i = 1, ..., r$, $f_i(\alpha_j) > 0$ $(j = 1, 2, 3)$ and there exists $i_0 \in \{1, ..., r\}$ such that $\alpha_4(f_{i_0}) > 0$; which is impossible because F is a fan.

Conversely, suppose S to be not basic open. If X is compact and non singular, by 2.8, 3.5 and 3.6 we have done. If not, take a birational model X_1 of X obtained compactifying and desingularizing X. Let S_1 be the strict transform of S in X_1. Then S_1 is not basic open and $\partial_z(S_1) \cap S_1 = \emptyset$, because S verifies these properties. So, by 2.8, 3.5 and 3.6, we can find a fan F of $\mathcal{K}(X_1)$ associated to a real prime divisor such that $\#(F \cap \tilde{S}_1) = 3$. Since $\mathcal{K}(X)$ and $\mathcal{K}(X_1)$ are isomorphic, F gives a fan G of $\mathcal{K}(X)$ such that G is associated to a real prime divisor and $\#(G \cap \tilde{S}) = 3$. □

4 The algorithms

By 2.3 (see also [ABF]) there is an algorithmic method for checking properties *(a)* and *(b)* of 2.8; so we can decide algorithmically if a semialgebraic S with $\partial_z S \cap S = \emptyset$ is open basic. This method works as follows:

1) It calculates $\dim(S^* \cap \partial_z S^*)$ by tecniques of cilindrical algebraic descomposition (C.A.D.), for instance (see [BCR]). If it is equal to 1, we know that S is not basic. If not,

we continue with 2).

2) It decides if some irreducible component of the exceptional divisor of the standard resolution of $\partial_z S$ is positive type changing using $\boxed{\text{A1}}$ and $\boxed{\text{A2}}$ in the points of ∂S which are not normal crossing (remark that $\boxed{\text{A2}}$ decides if the non-local completability at a point is due to a positive or a negative type changing component after some blowing-up).

Moreover, from [Vz] we have a complete description for fans in $\mathbb{R}(x,y)$ associated to a real prime divisor.

Consider the field $\mathbb{R}((u,v))$ of formal series in two variables over $\mathbb{R}$, with the ordering which extends 0^+ in $\mathbb{R}((u))$ by $v > 0$. Any ordering in $\mathbb{R}(x,y)$ is defined by an ordered $\mathbb{R}$-homomorphism $\psi : \mathbb{R}(x,y) \to \mathbb{R}((u,v))$ (see [AGR]). So a non-trivial fan F is given by 4 homomorphisms $\psi_1, \psi_2, \psi_3, \psi_4$. More precisely:

Theorem 4.1 (See [Vz]) *Let F be a fan in $\mathbb{R}(x,y)$. Then F is described as follows:*

1) *If F has as center an irreducible curve $H \subset \mathbb{S}^2$ and if $P(x,y) \in \mathbb{R}[x,y]$ is a polynomial generating the ideal $\mathcal{J}(H) \subset \mathbb{R}[x,y]$ of the image of H by a suitable stereographic projection, then*

$$\psi_i : \mathbb{R}(x,y) \to \mathbb{R}((t,z)),\ for\ i = 1,2,3,4$$

are defined (possibly interchanging x and y) as follows:

$$\begin{cases} \psi_1(x) = a_1 + \delta t^N \\ \psi_1(y) = a_2 + \sum_{i\geq 1} c_i t^{n_i} + z \end{cases} \qquad \begin{cases} \psi_2(x) = b_1 + \delta' t^M \\ \psi_2(y) = b_2 + \sum_{i\geq 1} d_i t^{m_i} + z \end{cases}$$

$$\begin{cases} \psi_3(x) = a_1 + \delta t^N \\ \psi_3(y) = a_2 + \sum_{i\geq 1} c_i t^{n_i} - z \end{cases} \qquad \begin{cases} \psi_4(x) = b_1 + \delta' t^M \\ \psi_4(y) = b_2 + \sum_{i\geq 1} d_i t^{m_i} - z \end{cases}$$

where $(a_1 + \delta t^N,\ a_2 + \sum_{i\geq 1} c_i t^{n_i})$ and $(b_1 + \delta' t^M,\ b_2 + \sum_{i\geq 1} d_i t^{m_i})$ are irreducible Puiseux parametrizations of two half-branches of H, centered respectively at (a_1, a_2), (b_1, b_2).

2) *If F is centered at a point $p \in \mathbb{S}^2$, we may suppose $p = (0,0)$ in a suitable stereographic projection, then*

$$\psi_i : \mathbb{R}(x,y) \to \mathbb{R}((z,t)),\ for\ i = 1,2,3,4$$

are given by one of the following expressions:

a)

$$\begin{cases} \psi_1(x) = t \\ \psi_1(y) = tw \end{cases} \qquad \begin{cases} \psi_2(x) = tw' \\ \psi_2(y) = t \end{cases}$$

$$\begin{cases} \psi_3(x) = -t \\ \psi_3(y) = -tw \end{cases} \qquad \begin{cases} \psi_4(x) = -tw' \\ \psi_4(y) = -t \end{cases}$$

with $w \in \{z + a, -z + a : a \in \mathbb{R}\}$, $w' \in \{z, -z\}$.

b) *Up to interchanging x and y,*

$$\begin{cases} \psi_1(x) = \delta t^N \\ \psi_1(y) = \sum_{i=1}^{s} c_i t^{n_i} + t^m w \end{cases} \qquad \begin{cases} \psi_2(x) = \delta t^N \\ \psi_2(y) = \sum_{i=1}^{s} c_i t^{n_i} + t^m w' \end{cases}$$

$$\begin{cases} \psi_3(x) = (-1)^N \delta t^N \\ \psi_3(y) = \sum_{i=1}^{s} (-1)^{n_i} c_i t^{n_i} + (-1)^m t^m w \end{cases} \qquad \begin{cases} \psi_4(x) = (-1)^N \delta t^N \\ \psi_4(y) = \sum_{i=1}^{s} (-1)^{n_i} c_i t^{n_i} + (-1)^m t^m w' \end{cases}$$

with $\delta \in \{1, -1\}$*;* $c_i \in \mathbb{R}$ *for* $i = 1, ..., s$*;* $N \leq n_1 < n_2 < \cdots < n_s$, g.c.d.$(N, n_1, ..., n_s) = d$ *and* g.c.d$(d, m) = 1$*; and* $w, w' \in \{z+a, -z+a, 1/z, -1/z : a \in \mathbb{R}\}$*, with* $w \neq w'$ *if* d *is odd and* $w \neq w', w \neq -w'$ *if* d *is even. If* $N = 1$*,* $c_1 = \cdots = c_s = 0$*, then* $w, w' \notin \{1/z, -1/z\}$*.*

c) *Up to interchanging* x *and* y*,*

$$\begin{cases} \psi_1(x) = \delta t^N \\ \psi_1(y) = \sum_{i=1}^{s} c_i t^{n_i} + t^m w \end{cases} \qquad \begin{cases} \psi_2(x) = (-1)^{N/d} \delta t^N \\ \psi_2(y) = \sum_{i=1}^{s} (-1)^{n_i/d} c_i t^{n_i} + t^m w' \end{cases}$$

$$\begin{cases} \psi_3(x) = \delta t^N \\ \psi_3(y) = \sum_{i=1}^{s} c_i t^{n_i} - t^m w \end{cases} \qquad \begin{cases} \psi_4(x) = (-1)^{N/d} \delta t^N \\ \psi_4(y) = \sum_{i=1}^{s} (-1)^{n_i/d} c_i t^{n_i} - t^m w' \end{cases}$$

with $\delta, c_i, N, n_i, w, w'$ *as in* b)*, but* d *always even and without supplementary conditions on* w, w'*.*

Remark 4.2 To each fan F in $\mathbb{R}(x, y)$ centered at $(0, 0)$ we can associate two families of arcs through $(0, 0)$ which are parametrized by $z \in (0, \epsilon)$. In fact, for each fixed $z \in (0, \epsilon)$, ψ_1 and ψ_3 (resp. ψ_2 and ψ_4) define the two half-branches of the same curve germ γ_1^z (resp. γ_2^z). This curves, γ_1^z and γ_2^z, verify one with respect to the other conditions a) and b) of 2.3 (see [ABF, 2.19]).

We want to know the relations between these two families of arcs associated to F and the families of arcs of $\boxed{\text{A1}}$ and $\boxed{\text{A2}}$ (2.3).

Let C be a curve germ through $(0, 0) \in \mathbb{R}^2$; and consider the standard resolution $\pi = \pi_N \cdots \pi_1$ of C at $(0, 0)$. For each $i = 1, \ldots, N$, denote by C_i the curve $(\pi_i \cdots \pi_1)^{-1}(C)$, by D_i the exceptional divisor arising during the i^{th} blowing-up, and by E_i the exceptional curve after i blowings-up (i.e. $E_1 = (\pi_i \cdots \pi_1)^{-1}(0, 0)$). D_i is an irreducible component of E_i.

Definition 4.3 *Let* F *be a fan of* $\mathbb{R}(x, y)$ *with center* $(0, 0) \in \mathbb{R}^2$*, for* $i = 1, ..., N$ *denote by* F_i *the fan obtained from* F *after* i *blowings-up by lifting the orderings of* F*. We say that* F *has the* property $\star(\rho)$ *with respect to* C *if it verifies:*

a) $F_{\rho-1}$ *is centered at the point* $0 = C_{\rho-1} \cap D_{\rho-1}$*.*

b) $F_{\rho-1}$ *is described in the sense of 4.1 as in* 2-a) *or* 2-b) *with* $N = 1$*,* $c_1 = \cdots = c_s = 0$*.*

c) $C_{\rho-1}$ *is not tangent to any curve of the two families associates to* $F_{\rho-1}$*.*

Remark 4.4 A fan F verifies $\star(\rho)$ with respect to a curve C if and only if for each $z \in (0, \epsilon)$ the curves γ_1^z and γ_2^z verify a) and b) of 2.3 with respect to C (see also [ABF, 2.9]).

Finally we have:

Theorem 4.5 *Let S be an open semialgebraic set in X, such that $\partial_z S \cap S = \emptyset$. Suppose that $\partial_z S^* \cap S^*$ is a finite set and S is not basic. Then, there exists an algorithmic method for finding a fan F of $\mathcal{K}(X)$ with $\#(F \cap \tilde{S}) = 3$.*

Proof: By 2.5 $\partial_z S$ has not type changing components with respect to σ_i^S; then by 2.8 there is at least one irreducible component D_ρ of the exceptional divisor of a standard resolution π of $\partial_z S$ at a point $O = (0,0)$ which is positive type changing with respect to some $\sigma'_i = \sigma_i^S \cdot \pi$. By 2.3 we can find algorithmically two arcs γ_1, γ_2 with the properties a) and b) of 2.3 with respect to an irreducible component of $\partial_z S$ through $p = \pi(D)$, for $\rho > 0$, such that γ_1 joins two regions in $(\sigma_i^S)^{-1}(1)$, while γ_2 joins a region in $(\sigma_i^S)^{-1}(1)$ to a region in $(\sigma_i^S)^{-1}(-1)$. Moreover, each γ_i $(i = 1,2)$ is defined by open conditions.

Then $(\gamma_1)_\rho$ and $(\gamma_2)_\rho$ are smooth arcs wich meet D_ρ transversally in different points $p, q \in D_\rho$ and ρ is the first level in the resolution process at wich γ_1 and γ_2 are separated. By [ABF, 2.19], γ_1 and γ_2 are parametrized by

$$\gamma_1 : \begin{cases} x = \delta t^N \\ y = \sum_{i=1}^s c_i t^{n_i} + f(t) \end{cases} \qquad \gamma_2 : \begin{cases} x = \delta' t^N \\ y = \sum_{i=1}^s d_i t^{n_i} + g(t) \end{cases}$$

where $\delta, \delta' \in \{1,-1\}$, $c_i, d_i \in \mathbb{R}$ are determined by [ABF, 2.19] as follows: $\delta' = \delta$ and $d_i = c_i$ or $\delta' = (-1)^{N/d}\delta$ and $d_i = (-1)^{n_i/d} c_i$, for $i = 1, ..., s$, with $d = \text{g.c.d.}(N, n_1, \dots, n_s)$; and $f(t) = at^m + ...$, $g(t) = bt^m + ...$ with $a \neq b$.

We can construct four fans with the property $\star(\rho)$ with respect to γ_1 and γ_2 as follows:

At the level ρ of the resolution process we have four fans centered at D_ρ in half-branches at p and q (Fig. 4.5), obtained by taking respectively a half-branch of D_ρ at p and an other at q.



Figure 4.5

Going back by the birational morphism $\pi_\rho \cdots \pi_1$ we obtain four fans centered at $O = (0,0)$ such that all they have the property $\star(\rho)$ with respect to γ_1 and γ_2. More precisely applying again [ABF, 2.19] we can describe them in terms of 4.1: for every pair $\eta, \eta' \in \{1,-1\}$ we have one of this fans $F_{\eta,\eta'}$ and his associated arcs are

$$\gamma_1^z : \begin{cases} x = \delta t^N \\ y = \sum_{i=1}^s c_i t^{n_i} + (\eta z + a) t^m \end{cases} \qquad \gamma_2^z : \begin{cases} x = \delta' t^N \\ y = \sum_{i=1}^s d_i t^{n_i} + (\eta' z + b) t^m \end{cases}$$

By the construction of γ_1, γ_2 it is easy to check that $\#(F_{\eta,\eta'} \cap \tilde{S}) = 3$. $\square$

Remark 4.6 In the hypotesis of 3.13 there are in fact infinite fans F of $\mathcal{K}(X)$ verifying $\#(F \cap \tilde{S}) = 3$, because there are infinite pairs of arcs joining respectively two region with sign 1 and a region with sign 1 with a region with sign -1.

So we find only fans with $w, w' \in \{z + a, -z + a : a \in \mathbb{R}\}$, according to description 4.1.2. In other case, $(\gamma_1^z)_\rho$ or $(\gamma_2^z)_\rho$ for some ρ, would be tangent to D_ρ, but applying $\boxed{\text{A1}}$ and $\boxed{\text{A2}}$ as in [ABF] we take γ_1, γ_2 without this property. This means that if it exists a fan F with w or $w' \in \{1/z, -1/z\}$ and $\#(F \cap \tilde{S}) = 3$, there is another fan F' with $w, w' \notin \{1/z, -1/z\}$ and $\#(F' \cap \tilde{S}) = 3$.

5 Principal sets

Using the results of the previous sections, we obtain a simple characterization of principal open (resp. closed) sets. In order to conserve the unity of this paper we give all results about principal sets in dimension 2, but in fact they can be extended to arbitrary dimension following similar proofs (Remarks 5.9). Details can be found in [Vz].

Let X be a compact, non singular, real algebraic surface.

Definition 5.1 *A semialgebraic set $S \subset X$ is* principal open *(resp.* principal closed*) if there exists $f \in \mathcal{R}(X)$ such that*

$$S = \{x \in X : f(x) > 0\}$$

$$(\textit{resp. } S = \{x \in X : f(x) \geq 0\})$$

Definition 5.2 *A semialgebraic set $S \in X$ is* generically principal *if there exists a Zariski closed set $C \in X$ with* $\dim(C) \leq 1$ *such that $S \setminus C$ is principal open.*

Remark 5.3 A semialgebraic set S is principal closed if and only if $X \setminus S$ is principal open.

Then it suffices to work with principal open sets.

Notation 5.4 Let S be an open semialgebraic set and $Y = \partial_{\mathrm{z}} S$. We denote by S^c the open semialgebraic set $X \setminus (S \cup Y)$ and by $A_1, \ldots, A_t$ (resp. $B_1, \ldots, B_l$) the connected component of S^c (resp. $S \setminus Y$).

Let σ_i be the sign distributions σ_i^S for $i = 1, \dots, t$ and σ_j^c be the sign distributions $\sigma_j^{S^c}$ for $j = 1, \dots, l$ defined as in 2.4. And denote by δ the total sign distribution defined by

$$\begin{aligned} \delta^{-1}(1) &= S \setminus Y \\ \delta^{-1}(-1) &= S^c \end{aligned}$$

Remark 5.5 A semialgebraic set S such that $\partial_{\mathrm{z}} S \cap S = \emptyset$ is principal open if and only if the sign distribution δ is admissible (that is, there exists $f \in \mathcal{R}(X)$ such that f induces δ on $X \setminus \partial_{\mathrm{z}} S$).

Theorem 5.6 *Let S be a semialgebraic set such that $\partial_{\mathrm{z}} S \cap S = \emptyset$. Then S is principal open if and only if $S^* \cap \partial_{\mathrm{z}} S^*$ and $(S^c)^* \cap \partial_{\mathrm{z}}(S^c)^*$ are finite sets.*

Proof: Suppose that S is principal then S and S^c are basic and by 2.8 no irreducible component of $\partial_{\mathrm{z}} S$ is positive type changing with respect to σ_i and σ_j^c for each $i = 1, ..., t$, $j = 1, ..., l$. Applying now 2.5 we have that $S^* \cap \partial_{\mathrm{z}} S^*$ and $(S^c)^* \cap \partial_{\mathrm{z}}(S^c)^*$ are finite sets.

Conversely suppose that $S^* \cap \partial_{\mathrm{z}} S^*$ and $(S^c)^* \cap \partial_{\mathrm{z}}(S^c)^*$ are finite sets, then by 2.5 again no irreducible component of $\partial_{\mathrm{z}} S$ is positive type changing with respect to σ_i and σ_j^c for $i = 1, ..., t$, $j = 1, ..., l$.

Remark that an irreducible component H of $\partial_{\mathrm{z}} S$ is positive (resp. negative) type changing with respect to δ if and only if H is positive type changing with respect to σ_i for some i (resp. σ_j^c for some j).

Hence no irreducible component of $\partial_{\mathrm{z}} S$ is type changing with respect to δ. Denote by Z^c the union of all the change components of $\partial_{\mathrm{z}} S$ and by Z the set of points where Z^c has dimension 1. So $[\underline{Z}] = [\underline{Z^c}]$ and $[Z] = 0$ in $\mathrm{H}_1(M, \mathbb{Z}_2)$, because it bounds the open sets $\mathrm{Int}(\overline{\sigma^{-1}(1)})$ and $\mathrm{Int}(\overline{\sigma^{-1}(-1)})$. Then by [BCR, 12.4.6] the ideal $\mathcal{J}(Z^c)$ of Z^c is principal. Let f be a generator of $\mathcal{J}(Z^c)$. Again by [BCR, 12.4.6] for each irreducible component H_k of $\partial_{\mathrm{z}} S$ not lying in Z^c we can choose a generator h_k of $\mathcal{J}(H_k)^2$ (wich exists because $2[H_k] = 0$). Then the regular function $f \cdot \prod h_k$ induces δ or $-\delta$. So δ is admisible and S principal. □

Remark that this proof is almost the same as the proof of [AB, Proposition 2].

Theorem 5.7 *Let S be a semialgebraic set in X.*

(1) S is principal open if and only if $\partial_{\mathrm{z}} S \cap S = \emptyset$ and for each fan F centered at a curve, $\#(F \cap \tilde{S}) \neq 1, 3$.

(2) S is principal closed if and only if $\partial_{\mathrm{z}} S \cap (X \setminus S) = \emptyset$ and for each fan F centered at a curve, $\#(F \cap \tilde{S}) \neq 1, 3$.

Proof: It is immediately using 5.6 and 3.5. □

Remarks 5.8 (1) Results 5.6 and 5.7 can be generalized to an arbitrary surface compactifying and desingularizing as in 2.9.

(2) All this section can be generalized to a compact, non singular, real algebraic set X, because the results of the previus sections used here (specifically 1.3 and 2.5) can be generalized to arbitrary dimension. Moreover, defining fan centered at a hypersurface H of X as a fan F associated to a real prime divisor V such that the prime ideal $\mathfrak{p} = \mathfrak{m}_V \cap \mathcal{R}(X)$ has height 1 and $\mathcal{Z}(\mathfrak{p}) = H$, we find an improvement of 4-elements fans criterion [Br, 5.3].

(3) For a compact, non singular, real algebraic set we obtain:
A semialgebraic set S is principal open (resp. closed) if and only if $\partial_z S \cap S = \emptyset$ (resp. $\partial_z S \cap (X \setminus S) = \emptyset$) and S is generically principal.

References

[AB] F. Acquistapace, F. Broglia: More about signatures and approximation, *to appear in Geometriae Dedicata.*

[ABF] F. Acquistapace, F. Broglia, E. Fortuna: When is a distribution of signs locally completable?, *to appear in Can. J. of Math.*

[AGR] M.E. Alonso, J.M. Gamboa, J.M. Ruiz: On orderings in Real Surfaces, *J. of Pure and Appl. Alg. 36 (1985) 1-14.*

[AR1] C. Andradas, J.M. Ruiz: More on basic semialgebraic sets, *in Real Algebraic and Analytic Geometry, 128-139, L.N.M. 1524, Springer-Verlag, Berlin-Heidelberg-New York 1992.*

[AR2] C. Andradas, J.M. Ruiz: On the ubiquity of Łojasewicz example, *to appear in Michigan Math. J.*

[BCR] J. Bocknak, M. Coste, M.F. Roy: Géométrie algébrique réelle, *Erg. Math. 12, Springer-Verlag, Berlin-Heidelberg-New York 1987.*

[BK] E. Brieskorn, H. Knörrer: Plane algebraic curves, *Basel Boston, Birkhäuser-Verlag, Stuttgart 1986.*

[Br] L. Brocker: On basic semialgebraic sets, *Exp. Math. 9 (1991) 289-334.*

[EC] F. Enriques, O. Chisini: Lezioni sulla teoria geometrica delle equazioni e delle funzioni algebriche, *3 Vols., Bologna 1915, 1918, 1924.*

[Rz] J.M. Ruiz: A note on a separation problem, *Arch. Math., Vol. 43, 422-426 (1984).*

[Vz] M.P. Vélez: La geometría de los abanicos en dimension 2, *to appear.*